

Aquatic insects in British Columbia: 100 years of study

**GEOFFREY G.E. SCUDDER, KAREN M. NEEDHAM,
REX D. KENNER**

**SPENCER ENTOMOLOGICAL MUSEUM, DEPARTMENT OF ZOOLOGY,
UNIVERSITY OF BRITISH COLUMBIA, VANCOUVER, BC, CANADA V6T 1Z4**

ROBERT A. CANNINGS

**ROYAL BRITISH COLUMBIA MUSEUM,
675 BELLEVILLE STREET, VICTORIA, BC, CANADA V8W 9W2**

and SYDNEY G. CANNINGS

**BRITISH COLUMBIA CONSERVATION DATA CENTRE,
MINISTRY OF SUSTAINABLE RESOURCE MANAGEMENT,
PO BOX 9344 STN PROV GOVT, VICTORIA, BC, CANADA V8W 9M1**

INTRODUCTION

Aquatic habitats in British Columbia are diverse and numerous. BC is home to large rivers, small ponds, and lakes which vary tremendously in size, depth, and chemical composition (Northcote and Larkin 1963), as well as a variety of fens, marshes, and bogs (Rosenberg and Danks 1987). Not surprisingly, studies in aquatic insects have a long history in our Province. The status of entomological knowledge from 1901 to 1951 in BC was summarized by Spencer (1952), including short paragraphs on each of the aquatic orders. The history of many of the entomologists involved and their collections are highlighted in Hatch's (1949) charming summary of a century of entomology (1835-1948) in the Pacific Northwest.

In the past, provincial studies have emphasized the importance of freshwater fisheries, and these have been accompanied by benthic invertebrate surveys. In recent years, forestry-fisheries interactions have come to the fore. The role of insects in this interaction is creating a new interest in aquatic entomology. The current concerns over water quality and groundwater pollution will undoubtedly involve more study of aquatic insects as environmental indicators.

EPHEMEROPTERA

Since the outstanding and prolific work of J. McDunnough in the 1920s and 1930s on the mayflies of Canada, the only BC-wide survey of Ephemeroptera was conducted by Filmer (1964). Most records since then have been obtained incidentally during other faunal studies. A checklist of BC species was produced by Scudder (1976). McCafferty and Randolph (1998) published a Canadian mayfly compendium that added seven new species to Scudder's 1976 list, as well as containing two species that Scudder (1976) omitted as probable erroneous records, for a total of 92 species in BC. Since this new list was compiled without examining specimens from the Royal British Columbia Museum, the Spencer Entomological Museum, or the Fisheries Branch of the Ministry of Water, Land and Air Protection, these collections may contain more new records for the Province.

Locally, Wigle and Thommasen (1990) collected and recorded ecological notes on 26 species of mayflies in the Bella Coola and Owikeno Lake watersheds of mid-coastal BC.

One of these, *Epeorus nitidus* Eaton, was a new record for the Province. Another new record for BC, *Acentrella turbida* (McDunnough), was found there later (McCafferty *et al.* 1994). Zloty (1996) and Zloty and Pritchard (1997) collected at various southwestern BC locales in search of *Ameletus* species, and to date have found one, *Ameletus pritchardi* Zloty, new to the Province and to science. S. Salter (pers. comm.) recently collected a new mayfly species for the Province, *Caenis youngi* Roemhild, during some invertebrate surveys at Liard Hot Springs in northern BC. In their comparison of the macroinvertebrate assemblages of coastal and continental streams and large rivers in southwestern British Columbia, Reece and Richardson (2000) found that *Paraleptophlebia temporalis* McDunnough predominated in coastal streams.

Heise (no date) in a study of the effect of logging on the fauna of streams in the Pentiction Creek and Sicamous Creek watersheds found that mayflies were very sensitive to logging. Richardson and Kiffney (2000) studied the response to metal pollutants of macroinvertebrate fauna in experimental streams off Mayfly Creek in the UBC Research Forest near Maple Ridge, and found that *Ameletus*, *Baetis*, and *Paraleptophlebia* (mainly *P. temporalis*) were very sensitive to Cu, Zn, Mn, and Pb, typical components of urban runoff. Rempel *et al.* (2000) in a study of the macroinvertebrates along gradients of hydraulic and sedimentary conditions in a 10-km reach of the Fraser River near Agassiz, found that the distribution of several genera of mayflies was correlated with hydraulic conditions. Rempel *et al.* (1999) also showed that the densities of *Baetis* and *Rhithrogena* were highest at a depth of 1.5 m before flooding, but shifted to depths of 0.2-0.5 m at peak flow.

Keys to genera of BC mayflies can be found in Edmunds *et al.* (1976), Merritt and Cummins (1996), and Needham (1996). Keys to species have only been completed for a few genera (e.g. *Ameletus*, Zloty (1996), Zloty and Pritchard (1997); *Caenis*, Provonsha (1990); *Tricorythodes*, Alba-Tercedor and Flannagan (1995)).

ODONATA

Spencer (1952) in his brief summary of the status of the knowledge of Odonata in British Columbia did not cite what is perhaps the most significant work of the period, Whitehouse's (1941) delightful and detailed treatment of the provincial fauna. He also neglected Walker's important monographs on two of our most speciose genera, *Aeshna* (Walker 1912) and *Somatochlora* (Walker 1925); although these dealt with the entire North American fauna of these genera, they included much information on the species in British Columbia.

Walker's major opus on the Odonata of Canada and Alaska (Walker 1953, 1958) led off the second half of the century and was completed by the international Odonata expert P. Corbet (Walker and Corbet 1975). It listed many of the locality records known to this time and summarized ecological information.

Since the mid 1970s the study of the Odonata in BC has been continuous. It has been helped by work in adjacent areas, particularly the Biological Survey of Canada's Yukon insect project (S.G. Cannings and R.A. Cannings 1997; S.G. Cannings *et al.* 1991) and the extensive studies of Paulson (e.g. 1997, 1999) to the south in Washington State. This followed the annotated checklist produced by Scudder *et al.* (1976) and the book written by R.A. Cannings and Stuart (1977). The latter published the first distribution maps for the species of the Province.

Robert and Sydney Cannings, periodically helped by their brother Richard, have been the driving force in odonatology in the Province during the past 25 years. They organized numerous dragonfly inventories across the Province, from the Brooks Peninsula on the outer coast of Vancouver Island (R.A. Cannings and S.G. Cannings 1983), to the grasslands of the Chilcotin (R.A. Cannings and S.G. Cannings 1987), to the Rocky



F.C. Whitehouse, a pioneering odonatologist, collecting specimens from his caravan at Harrison and Cultus Lakes in the summer of 1936.

Mountains (R.A. Cannings *et al.* 2000) and the North (R.A. Cannings *et al.* 2001). Others who have made significant contributions to these collections and inventories include G. Hutchings, L. Ramsay, C. Guppy, R. Kenner, D. Blades, and H. Nadel. In addition, the Cannings, along with their colleagues, have documented new species to the Province (R.A. Cannings 1988, 1997; Kenner 2000a) and published much faunistic (R.A. Cannings 1978, 1996; R.A. Cannings *et al.* 1980; S.G. Cannings 1980; S.G. Cannings and R.A. Cannings 1994) and ecological work (R.A. Cannings 1980, 1982a; R.A. Cannings *et al.* 1980; Paulson and R.A. Cannings 1980). Significant in the taxonomic work done on the Odonata in British Columbia and Yukon over the past 25 years is the description of four previously unknown larvae (R.A. Cannings 1981; S.G. Cannings and R.A. Cannings 1980; R.A. Cannings and Doerksen 1979; Kenner *et al.* 2000) and the redescription of several others, along with improved keys to this important life stage (R.A. Cannings 1982b; Kenner 2001).

G.P. Doerksen had begun to examine local dragonflies intensively (Doerksen 1980) when he was killed by a Grizzly Bear while on a research trip to Liard River Hot Springs in 1981. His wonderful dragonfly photographs, willed to the Royal BC Museum, are a valuable research and interpretative resource. Peters (1998), while on a visit from Germany, published a useful study on variability in *Aeshna*. R. Kenner (Kenner 1996, 2000b, 2000c; Kenner and Lane 1997; Kenner and R.A. Cannings 2001) and I. Lane (Lane 2000) have improved our knowledge of the fauna of the Lower Mainland, including the documentation of the first Canadian breeding site of *Tanypteryx hageni* Selys.

Few experimental studies on the Odonata have been undertaken in the Province. G. Pritchard, University of Calgary, has studied the ecology and development of Odonata, especially *Argia vivida* Hagen, of warm springs in British Columbia and adjacent regions (Pritchard 1989, 1991). At the University of BC, P. Pearlstone (1973) examined the food

of larval *Enallagma boreale* Selys and R. Baker (1983) studied the larval competition and feeding behaviour of *Ischnura cervula* Selys.

An important recent development has been the listing of species of management concern by the BC Conservation Data Centre (Ramsay and S.G. Cannings 2000; R.A. Cannings 2001). At the end of the 2001 field season, 23 species were listed. This highlighting of rare species spurred research into their status and habitats, and stimulated regional inventories.

Odonata are now the best known aquatic insects in the Province. Of course, there is still much to learn, especially regarding the ecology of the fauna and the distribution of northern species. Spencer noted 78 species in 1952, but the most accurate published record up to that time was from Whitehouse (1941), who recognized 74 species in 23 genera. By 1977, R.A. Cannings and Stuart could report 80 species and 23 genera. In 2001, 87 species in 27 genera are known from the Province (there are 201 recorded in Canada). Perhaps the most striking addition to the provincial list in the past 50 years was the discovery of *Calopteryx aequalis* Say at Christina Creek in 1998 (R.A. Cannings *et al.* 2000). This added a new family to British Columbia, the Calopterygidae, making a total of 10.

All available records for the Odonata in BC are databased, georeferenced, and mapped, ready to serve as baseline data for the faunal changes of the new millennium. Useful works for identification, in addition to Walker (1953, 1958), Walker and Corbet (1975), and R.A. Cannings and Stuart (1977) are Westfall and May (1996), Needham *et al.* (2000), and Dunkle (2000).

PLECOPTERA

Ricker (e.g. 1943, 1954) conducted early systematic and distributional studies on stoneflies. An annotated checklist of the Plecoptera of BC was published by Ricker and Scudder (1976). S.G. Cannings (1989) also provided a number of new records, especially for capniids. An up-to-date checklist can be extracted from the North American stonefly list provided on the internet by Stark (2001). Currently, 132 species are known from the Province. More detailed systematic studies that are relevant include Nelson and Baumann's (1989) work on *Capnia*, Stanger and Baumann's (1993) work on *Taenionema*, and Stark and Nelson's (1994) work on *Yoraperla*. Kenneth Stewart (University of North Texas) and Mark Oswood are actively writing "Stoneflies of Alaska and Northwestern Canada," which will include history, keys, illustrations, new locality records, maps, species accounts, and biological notes on the species documented for Alaska, British Columbia, Yukon, and Northwest Territories; publication is anticipated in 2003.

D.B. Donald and R.S. Anderson of the Canadian Wildlife Service (Edmonton) surveyed the stoneflies of the Rocky Mountain National Parks. The lentic stoneflies of these collections were summarized in Donald and Anderson (1980). Donald and Patriquin (1983) related the wing length of Rocky Mountain lentic capniids to their inferred postglacial recolonization history.

A number of studies investigating stonefly population dynamics, foraging ecology, species interactions, and life history in coastal streams have been undertaken by J. Richardson and his colleagues at UBC (Reece and Richardson 2000; Rempel *et al.* 2000; Richardson 2002; Soluk and Richardson 1997). Muchow and Richardson (2000) reported the occurrence of Plecoptera in small headwater streams in the UBC Research Forest near Maple Ridge. They found the fauna differed in intermittent and continuous flow streams. While *Paraleuctra vershina* Gautin and Ricker and *Visoka cataractae* (Neave) occurred only in continuous flow streams, *Alloperla pilosa* Needham and Claassen, *Despaxia augusta* (Banks), *Moselia infusca* (Claassen), *Ostracercia foesteri* (Ricker), *Pteronarcys californica* Newport, *Soyedina producta* (Claassen), *Zapada cinctipes* (Banks), and *Z. oregonensis* (Claassen) occurred in both. Even in small streams (less than half a metre wide) with intermittent flow, the univoltine *Moselia infusca*, *Soyedina producta*, and

Zapada cinctipes were found to emerge in periods when no flow was perceptible. *Despaxia augusta*, with a 2-year life cycle, was also able to complete its development in the periodically 'dry' channels and reached its highest densities in intermittent streams. The authors suggest that suitable refugia exist for this species and others in the wet sediments of these habitats despite the periodic disappearance of detectable surface flows.

All available, reliably identified records are now databased, georeferenced, and mapped. References useful for identification include Baumann *et al.* (1977), Harper and Stewart (1984), Jewett (1959), Ricker (1943), Stark *et al.* (1986), and Stewart and Stark (1988).

HEMIPTERA

In addition to records in the early literature of Hemiptera in British Columbia, Scudder (1977) published an annotated checklist of the aquatic and semi-aquatic bugs in the Province. In the recent checklist for Canada (Maw *et al.* 2000), a total of 46 species of aquatics (Infraorder Nepomorpha: Families Belostomatidae, Nepidae, Gelastocoridae, Corixidae, Notonectidae) and 14 species of semi-aquatics (Infraorder Gerromorpha: Families Mesoveliidae, Hebridae, Hydrometridae, Gerridae, Veliidae) are recorded.

Over the past 40 years, detailed studies of the biology of both Corixidae and Gerridae have been undertaken, much of this associated with a study of the saline lakes of the British Columbia interior. Scudder (1965a, 1969a, 1969b) investigated the distribution of two closely related species of *Cenocorixa* in these lakes, and found that while *C. bifida* (Hung.) was evidently a freshwater species, *C. expleta* (Uhler) was more saline tolerant.

Extensive study of the osmotic and ionic balance in these two corixid species (Scudder 1971a; Scudder *et al.* 1972; Szibbo and Scudder 1979; Needham 1990) and an investigation of their cuticular permeability (Oloffs and Scudder 1966; S.G. Cannings 1981; S.G. Cannings *et al.* 1988) was accompanied by detailed histological and ultrastructural studies of their excretory organs (Jarial and Scudder 1970) and other organs that were evidently involved in osmotic and ionic regulation (Jarial *et al.* 1969; Lo and Acton 1969; Jarial and Scudder 1971). Although the two species differed in their osmotic and ionic regulatory abilities at high salinities, both were able to regulate equally well in low salinity waters. Indeed, the more saline tolerant *C. expleta*, which in the field does not live in freshwater lakes, was successfully reared in freshwater (S.G. Cannings 1978). These studies subsequently led to much more detailed research on the coastal *C. blaisdelli* (Hung.) (Cooper *et al.* 1987, 1988, 1989) and other corixid species (Scudder 1987).

Both of the interior species of *Cenocorixa*, and other Corixidae, were found to be predaceous (Jansson and Scudder 1972; Reynolds 1975), which permitted laboratory rearing and led to other studies. Once species had been reared and the immature stages of both *C. bifida* and *C. expleta* described (Scudder 1966a), studies were carried out on their life cycle (Jansson and Scudder 1974), flight muscle development (Scudder 1971b), and flight muscle polymorphism (Scudder 1964; Acton and Scudder 1969; Scudder and Meredith 1972; Scudder 1975).

Jansson (1972, 1973, 1974a, 1974b), in his extensive study of stridulation in these *Cenocorixa* species, found species-specific mating signals and hence an effective pre-mating isolating mechanism. Further, in an intensive study of their feeding biology, using serological and other techniques, Reynolds and Scudder (1987a, 1987b) studied both the fundamental and realized feeding niches of *C. bifida* and *C. expleta*.

In a review of the factors governing the distribution of *C. bifida* and *C. expleta*, Scudder (1983), following studies on mite parasitism of these species (Smith 1977), suggested that *C. expleta* was excluded from freshwater lakes by mite parasitism. This suggestion has been substantiated by more research (Bennett and Scudder 1998), which showed that for

C. expleta saline lakes are enemy-free space. This research has been cited as an example of a parasite-structured animal community (Minchella and Scott 1991).

In the Gerridae, several species were found to coexist in BC lakes (Maynard 1969; Scudder 1971c). Following the rearing and description of the immature stages of these (Scudder and Jamieson 1972; Spence and Scudder 1978), detailed studies were undertaken of their food consumption and predatory behaviour (Jamieson and Scudder 1977, 1979), growth patterns (Spence *et al.* 1980a), life cycles (Spence and Scudder 1980; Rowe and Scudder 1990), mating and other behaviours (Spence *et al.* 1980b; Rowe 1992, 1994). Spence (1981, 1983, 1989) was then able to undertake a detailed analysis of habitat selection, life cycle strategy, species-packing, and coexistence in these pond skaters. Subsequently, Spence (1990) discovered introgressive hybridization in two species of *Limnoporus* within British Columbia and Alberta, and has gone on to look at the mating system of these (Spence and Wilcox 1986; Wilcox and Spence 1986), and some of the genetic factors accompanying this hybridization between non-sister species (Spence and Maddison 1986; Sperling and Spence 1990, 1991; Sperling *et al.* 1997).

Seven species of Notonectidae occur in BC (Scudder 1965b, 1977). *Notonecta borealis* Hussey was discovered to be primarily a flightless species (Scudder 1966b). Ellis and Borden (1969) have investigated the effect of temperature and other environmental factors on *N. undulata* Say.

MEGALOPTERA

Spencer (1942) treated the Megaloptera, which includes the dobsonflies and the alderflies, within his listing of the Neuroptera, and recorded 4 species from British Columbia. A systematic review of the Sialidae by Ross (1937) gives a key to the species of *Sialis*; 5 species are now known to occur in the Province. Munroe (1951, 1953) sorted out the identity of material from British Columbia that Spencer (1942) reported as *Neohermes disjunctus*, and now 3 species of Corydalidae are known from BC.

While *Dysmicohermes disjunctus* (Walker) is widely distributed, both *Chauliodes pecticornis* L. and *Protochauliodes spenceri* Munroe may be at risk, as they have a restricted distribution in areas of the Province subject to heavy human impact. *Chauliodes pecticornis* is known from Cloverdale and Cowichan, while *Protochauliodes spenceri* is restricted to south-east Vancouver Island in Canada. All known records of Megaloptera in the Province are databased, georeferenced, and mapped.

NEUROPTERA

Of the 8 families of Neuroptera reported from British Columbia, only the family Sisyridae has aquatic larvae. In the Sisyridae, although Spencer (1942) recorded only *Sisyrha vicaria* (Walker) from BC, based upon a specimen from Agassiz determined by F.M. Carpenter, it is now known that another species of spongilla fly, *Sisyrha fuscata* (Fab.), also occurs in the Province. While *S. vicaria* is widely distributed, *S. fuscata* is only known from Kaslo and Seton Lake. A key to the species of Sisyridae is given in Parfin and Gurney (1956). All known records of aquatic Neuroptera in BC are now databased, georeferenced, and mapped.

TRICHOPTERA

Some of the earliest records of Trichoptera from British Columbia come from N. Banks, who worked on the group from the early 1900s to the early 1940s. H.H. Ross also contributed significantly to these early records, collecting in the 1930s, 1940s, and 1950s throughout North America. The first attempt to pull all of these scattered records together

into one publication was made by Ross and Spencer (1952), who created a preliminary list of the Trichoptera of BC. This list was added to significantly by Nimmo and Scudder (1978) when they published their annotated checklist containing 248 species, 43 of which were new records for BC. At this time, they stated that "there has been no concerted Trichoptera collecting in British Columbia as a whole, so the fauna must still be regarded as incompletely known."

To partially rectify this, A. Nimmo spent the spring, mid-summer, and fall of 1979 exploring much of the Province in search of caddisflies (Nimmo and Scudder 1983). He added many range extensions to known species, and also recorded 36 new species for BC (Nimmo and Scudder 1983), thus bringing the provincial total to 279 species (they removed 5 species from the 1978 list). However, Nimmo admitted that the northern half of the Province, the Queen Charlotte Islands, and many alpine areas still remain uncollected (Nimmo and Scudder 1983).

Also in the 1970s, G. Wiggins published his invaluable and beautifully illustrated identification guide to the larval caddisflies of North America (Wiggins 1977). A second edition in 1996 recognized some 1400 species of Trichoptera in North America by the end of 1993 (Wiggins 1996). More specific to our region are the keys provided by Schmid (1980) to the genera of Trichoptera in Canada and adjacent States. An English version of this Insects and Arachnids of Canada (Part 7) publication has also been recently made available (Schmid 1998). Unfortunately, keys to caddisflies of BC at the species level are still lacking.

Collections of caddisflies from specific areas of our Province can be found in Clemens *et al.* (1939) for Okanagan Lake, Hardy (1955) for the Forbidden Plateau area of Vancouver Island, Nimmo (1971, 1974, 1977) for eastern BC, Rawson (1934) for the Kamloops region, Schmid and Guppy (1952) for southern Vancouver Island, Scudder (1969b) for the Fraser Plateau, and Winterbourn (1971a,b) for Marion Lake, BC.

On the biology of caddisflies, Richardson (1991) found *Lepidostoma roafi* (Milne) to be univoltine. In a study contrasting the fauna of coastal and continental streams and large rivers in southwestern BC, Reece and Richardson (2000) found that *Onocosmoecus unicolor* (Banks) was unique to coastal streams, while *Brachycentrus occidentalis* Banks was unique to larger rivers. *B. americanus* (Banks) was unique to continental streams in the Merritt area. In Marion Lake, Trichoptera larvae, in particular *Banksiola crotchii* Banks, were the most important food item of rainbow trout (Efford and Tsumura 1973). The effects of the mosquito larvicide *Bacillus thuringiensis* on caddisflies was reported by Duckitt (1986).

Rempel *et al.* (1999), in an investigation of the effect of flooding on benthic invertebrates in the Fraser River near Agassiz, noted that *Hydropsyche* was most abundant at 1.5 m in all months of the years, but the location shifted laterally over a distance of 30 m through the flood cycle. They evidently migrated to the shore zone with the rise in water level, so maintaining a suitable hydraulic microhabitat. Heise (no date), in a study of the effect of logging on the fauna of streams in Penticton Creek and Sicamous Creek watersheds, found that while some caddisfly larvae showed a constant pattern of distribution in response to logging, *Psychoglypha*, a shredder, declined after logging and *Ecclisocosmoecus*, an algal feeder, increased after logging.

Caira and Scudder (1985) reported parasitism of *Psychoglypha alascensis* Banks (Trichoptera: Limnephilidae) by *Pseudoalloeceadium alloneotenicum* (Wootton) (Digenea: Alloeceadiidae). Caira (1981) investigated parasitism of Trichoptera by *Bunodera mediovittellata* Zimbaluk & Roytman (Digenea: Alloeceadiidae) and the encapsulation response. This hemocyte encapsulation response in *P. alascensis* was studied later using Epon implants (Caira and Scudder 1987).

COLEOPTERA

British Columbia has a very rich aquatic beetle fauna. For example, BC has more species of Dytiscidae, Amphizoidae, Haliplidae, and Hydraenidae than any other Province (Larson *et al.* 2000; Bousquet 1991). Yet, beetles appear to be one of the more neglected groups of aquatic insects in BC. Early lists of Coleoptera from various parts of the Province (e.g. Keen 1895, 1905; Stace-Smith 1929, 1930) do contain some aquatic species, but there has been no published provincial checklist, although a preliminary list was developed by Scudder (unpublished). Most of the works dealing with aquatic beetles in BC have been as parts of studies of particular taxa over larger geographical areas. References to keys for aquatic beetles are in Bousquet (1991) and Arnett and Thomas (2001). Species lists for BC are in Bousquet (1991) and Larson *et al.* (2000).

The most important person to study aquatic beetles in BC was H.B. Leech. Leech was born in Kamloops, received much of his early education in Vernon, and graduated from UBC in 1933. From 1930-1947, he worked at the Forest Entomology Laboratory in Vernon where he collaborated with R. Hopping. In 1947, he joined the California Academy of Sciences. He described his first new species in 1937, *Agabus vancouverensis* Leech (Leech 1937), and subsequently described over 50 beetle taxa (Kavanaugh and Arnaud 1981). His collection of over 30,000 specimens of aquatic beetles, many from BC, was donated to the California Academy of Sciences in 1947. Several species of aquatic beetles found in BC are named after Leech: *Haliplus leechi* Wallis, *Helophorus leechi* McCorkle, and *Cymbiodyta leechi* Miller, which is known from Washington and probably occurs in BC (Smetana 1988).

Several other people are associated with aquatic beetles in BC mostly as collectors; for example, G.J. Spencer, R. Hopping, and G. Stace-Smith. M.H. Hatch described several species of aquatic beetles found in BC and is best known for his "Beetles of the Pacific Northwest" (Hatch 1953-1971). Each of them has at least one aquatic beetle from BC named in his honor.

In more recent times, Scudder (1969b) reported some species from interior saline lakes, and he, his students, and associates have collected aquatic beetles as part of more general surveys and added to our knowledge of the distribution of various species, especially in the Gulf Islands and in the northern part of the Province. In addition, J. Lancaster has conducted an ecological study of aquatic beetles in a comparison of the community structure in a series of saline lakes on the Chilcotin Plateau (Lancaster and Scudder 1987). Kenner (2000d) described the distribution of two species of gyinids in BC, the Yukon, and Alaska.

DIPTERA

The Manual of Nearctic Diptera (McAlpine 1981, 1987) is a wonderful resource for information on systematics and biology of North American Diptera. Illustrated keys identify immature and adult specimens to family and genus. The taxonomic and distributional status, to the early 1960s, of many British Columbia species is outlined in Stone *et al.* (1965).

The Mountain Midges (Deuterophlebiidae) are peculiar, little-known flies found in streams in the western mountains. Courtney (1990) documented the systematics of the Nearctic species, including those known from BC.

In the Chaoboridae, Northcote (1964) and Teraguchi and Northcote (1966) studied the vertical distribution and vertical migration of *Chaoborus* larvae in BC lakes. Taxonomic works on the Phantom Midges that relate to the provincial fauna include Saether (1970) and Borkent (1979, 1993).

Early work on the Culicidae in BC includes that of Hearle (1926, 1927) and Curtis (1967). Belton (1978) studied the mosquitoes of Burnaby Lake, and more recently produced a monograph on the 46 species known in BC (Belton 1983). This work, together with Wood *et al.* (1979), can be used to identify these species.

In his study of the fauna of saline lakes on the Fraser Plateau, Scudder (1969b) found that *Aedes campestris* Dyas and Knab occurred in the most saline of them. The osmotic and ionic regulation in the larvae of this mosquito is therefore of interest; it has been studied by Phillips and Meredith (1969a). They also investigated the active transport of sodium and chloride by the anal papillae of the larvae (Phillips and Meredith 1969b) and the electron microscopic structure of these organs (Meredith and Phillips 1973a, 1973b). Regulation and active transport of magnesium by the Malpighian tubules of these larvae was described by Kiceniuk and Phillips (1974) and Phillips and Maddrell (1974). They also discovered that these tubules could actively transport sulphate ions (Maddrell and Phillips 1975). This work led to comparative studies on other saline-water mosquito larvae including *Aedes dorsalis* (Meigen) and *A. togoi* (Theobald) (Meredith and Phillips 1973c; Strange *et al.* 1982, 1984; Strange and Phillips 1984), both of which occur in the Province. Important reviews resulted from this research (Bradley and Phillips 1977; Phillips *et al.* 1978; Phillips and Bradley 1978; Strange and Phillips 1985; Ng and Phillips 1985).

Gammarid predation on *Aedes togoi* at Horseshoe Bay was reported by Hossack and Costello (1979). Oviposition of *Culex pipiens* L. in water at different temperatures was described by Gillespie and Belton (1980). Ishii and Belton (1984) provided evidence for autogenous egg development in this species and Belton (1982) described the cuticular vestiture of *Aedes communes* (DeGeer) and *A. nevadensis* Chapman and Burr.

The comprehensive treatment of the Black Flies (Simuliidae) of North America by P. Adler, D. Currie, and D.M. Wood is nearing completion; when published, this work will deal with all aspects of the systematics of BC species. Early studies by Curtis (1954) and Shewell (1957, 1959) were followed by research by Peterson (1970), Mahrt (1982), Currie and Adler (1986), and Corkum and Currie (1987). Currie's work in Alberta (1986) and the Yukon (1997) has direct application to the BC fauna. Studies on *Parasimulium* (Borkent 1992; Borkent and Currie 2001) have led to a better understanding of the phylogenetic origins of the Simuliidae. Currie and Walker (1992) documented the use of fossil black fly larvae in paleoecological studies. Cytological research into the giant chromosomes of simuliids has been undertaken in the genus *Prosimulium* (Basrur 1962), in *Stegopterna* (Madahar 1969), in the *Simulium* (*Eusimulium*) *vernum* group (Hunter and Connolly 1986), and in the *S. (E.) aureum* group (Leonhardt 1985).

McMullen (1978) surveyed Ceratopogonidae in the Okanagan area of BC. Light trap catches of *Culicoides* in the Fraser Valley were described by Costello (1982). Anderson and Belton (1993) examined populations of *Culicoides obsoletus* Meigen in the lower mainland of BC and hypersensitivity of horses to *Culicoides* bites was reported (Anderson *et al.* 1988, 1991, 1993, 1996). A. Borkent of Enderby is a world authority on Ceratopogonidae and some of his systematic studies directly relate to BC (Borkent and Grogan 1995; Borkent 1998).

Both freshwater and saline-tolerant species of the Chironomidae inhabit saline lakes in the BC interior (Scudder 1969b). At the University of Victoria, Morley and Ring (1972a, 1972b) studied the life histories, ecology, and identification of the intertidal chironomids *Paraclunio alaskensis* Coquillett, *Saundersia clavicornis* (Saunders), *S. marinus* (Saunders), and *S. pacificus* (Saunders). The detailed study of the occurrence of midges in the littoral zone (R.A. Cannings and Scudder 1978) resulted in the discovery of a new species of *Chironomus* (R.A. Cannings 1975a) and many species new to BC and Canada (R.A. Cannings 1975b). Subsequently, the physiology of this and other *Chironomus* species was investigated (Sargent 1978) and the phenology compared (R.A. Cannings and Scudder

1979). The ecology of one of these, *C. tentans* Fab., which has distinct chromosomal races in the Province (Acton and Scudder 1971), was researched by Topping (1971, 1972) and, experimentally, Topping (1969) and Topping and Acton (1976) examined the influence of environmental factors on the frequency of certain chromosomal inversions. Eastern Brook Trout were used as predators in this research. The chironomids of Marion Lake in the Fraser Valley were studied by Hamilton (1965); Schultze and Northcote (1972) documented chironomids as fish food in another coastal lake. Mundie (1971) studied their diel drift in Robertson Creek, an artificial spawning channel and secondary outflow of Great Central Lake on Vancouver Island, and found that larval drift rates increased almost two-fold in darkness.

Brillia retifinis Saether is a multivoltine species, able to track shifts in resource abundance in its habitat by virtue of its short generation time (Richardson 1991). The influence of *Brillia* on the decomposition of conifer leaves in a coastal stream was reported by Summerbell and R.A. Cannings (1981). Richardson and Kiffney (2000), in their study of the response of aquatic insects to metal pollutants, found that chironomids (mainly *B. retifinis*) showed no significant decrease in densities with increasing doses of copper, zinc, and manganese, typical urban pollutants. Parkinson and Ring (1983) examined more physiological adaptations of *P. alaskensis* to the marine environment. As part of a provincial government attempt to control the introduced aquatic weed, *Myriophyllum spicatum* L., in warm lakes and waterways, potential biological control agents were screened. The chironomids found in this study were documented by Kangasniemi and Oliver (1983). One of them, an unnamed, native species that feeds on the plant's apical buds, was subsequently described as *Cricotopus myriophylli* by Oliver (1984). Borkent's (1984) systematic work on the *Stenochironomus* complex relates to the BC fauna. Ian Walker and his students and colleagues (Walker *et al.* 1991, 1993; Heinrichs *et al.* 1999; Rück *et al.* 1998; Palmer 1998) have used chironomid fossils in lake sediments to reconstruct paleoenvironments in British Columbia. The larvae of Canadian chironomids can be identified to genus using Oliver and Roussel (1983).

CONCLUSION

The above clearly documents that there has been a dramatic increase in our knowledge of the aquatic insects in British Columbia over the past century. However, much remains to be done, with the Ephemeroptera in particular needing more attention. Additionally, many areas of the province remain to be surveyed, such as the far North and the numerous islands off the west coast. Many more interesting discoveries can be expected during the next 100 years.

ACKNOWLEDGEMENTS

The authors thank the following people for their contributions to this paper: Peter Belton, Art Borkent, Doug Currie, Tom Northcote, John Phillips, and John Richardson. Thanks also to Peter Belton for his thoughtful and meticulous review of the manuscript. His comments greatly improved our submission.

REFERENCES

- Acton, A.B. and G.G.E. Scudder 1969. The ultrastructure of the flight muscle polymorphism in *Cenorcorixa bifida* (Hung.) (Heteroptera, Corixidae). *Zeitschrift für Morphologie und Ökologie der Tiere* 65: 327-335.
- Acton, A.B. and G.G.E. Scudder. 1971. The zoogeography and races of *Chironomus* (= *Tendipes*) *tentans* Fab. *Limnologia* 8: 83-92.
- Alba-Tercedor, J. and J. Flannagan. 1995. Two new Canadian species of the genus *Tricorythodes* Ulmer, with additional studies on other North American species (Insecta, Ephemeroptera: Leptohyphidae). *Canadian Journal of Zoology* 73: 1588-1598.
- Anderson, G.S. and P. Belton. 1993. A population study of *Culicoides obsoletus* (Diptera: Ceratopogonidae) in the lower mainland of British Columbia. *The Canadian Entomologist* 125: 439-442.
- Anderson, G.S., P. Belton and N. Kleider. 1988. The hypersensitivity of horses to *Culicoides* bites in British Columbia. *Canadian Veterinary Journal* 29: 718-723.
- Anderson, G.S., P. Belton and N. Kleider. 1991. *Culicoides obsoletus* Meigen (Diptera: Ceratopogonidae) as a causal agent of *Culicoides* hypersensitivity (sweet itch) in British Columbia. *Journal of Medical Entomology* 28: 685-693.
- Anderson, G.S., P. Belton and N. Kleider. 1993. Hypersensitivity of horses in British Columbia to extracts of native and exotic species of *Culicoides* (Diptera: Ceratopogonidae). *Journal of Medical Entomology* 30: 657-663.
- Anderson, G.S., P. Belton, E. Jähren, H. Lange and N. Kleider. 1996. Immunotherapy trial for horses in British Columbia with *Culicoides* (Diptera: Ceratopogonidae) hypersensitivity. *Journal of Medical Entomology* 33: 458-466.
- Arnett, R.H. and M.C. Thomas. 2001. *American Beetles: Vols. 1 and 2*. CRC Press, New York.
- Baker, R.L. 1983. Spacing behaviour by larval *Ischnura cervula* Selys: effects of hunger, previous interactions, and familiarity with an area (Zygoptera: Coenagrionidae). *Odonatologica* 12: 201-207.
- Basrur, P.K. 1962. The salivary gland chromosomes of seven species of *Prosimulium* (Diptera: Simuliidae) from Alaska and British Columbia. *Canadian Journal of Zoology* 40: 1019-1033.
- Baumann, R.W., A.R. Gaufin and R.F. Surdick. 1977. The Stoneflies (Plecoptera) of the Rocky Mountains. *Memoirs of the American Entomological Society* 31: 1-208.
- Belton, P. 1978. The mosquitoes of Burnaby Lake, British Columbia. *Journal of the Entomological Society of British Columbia* 75: 20-22.
- Belton, P. 1982. The cuticular vestiture of larvae of *Aedes communis* and *A. nevadensis* (Diptera: Culicidae). *Canadian Journal of Zoology* 60: 1642-1646.
- Belton, P. 1983. The Mosquitoes of British Columbia. *British Columbia Provincial Museum Handbook* 41. 188 pp.
- Bennett, A.M.R. and G.G.E. Scudder. 1998. Differences in attachment of water mites on water boatmen: further evidence of differential parasitism and possible exclusion of a host from part of its potential range. *Canadian Journal of Zoology* 76: 824-834.
- Borkent, A. 1979. Systematics and bionomics of species of the subgenus *Schadonphasma* Dyar and Shannon (*Chaoborus*, Chaoboridae, Diptera). *Quaestiones Entomologicae* 15: 122-255.
- Borkent, A. 1984. The systematics and phylogeny of the *Stenochironomus* complex (*Xestochironomus*, *Harrisius* and *Stenochironomus*) (Diptera: Chironomidae). *Memoirs of the Entomological Society of Canada* 128: 1-269.
- Borkent, A. 1992. *Parasimulium* (Diptera: Simuliidae), a black fly genus new to the extant fauna of Canada. *The Canadian Entomologist* 124: 743-744.
- Borkent, A. 1993. A world catalogue of fossil and extant Corethrellidae and Chaoboridae (Diptera) with a listing of references to keys, bionomic information and descriptions of each known life stage. *Entomologica Scandinavica* 24: 1-24.
- Borkent, A. 1998. A revision of *Neurobezzia* Wirth & Ratanaworabhan and *Neurohelea* Kieffer, with a description of a new genus and discussion of their phylogenetic relationships (Diptera: Ceratopogonidae). *Entomologica Scandinavica* 29: 137-160.
- Borkent, A. and D.C. Currie. 2001. Discovery of the female of *Parasimulium* (*Astoneomyia*) *melanderi* Stone (Diptera: Simuliidae) in a cave in British Columbia, with a discussion of its phylogenetic position. *Proceedings of the Entomological Society of Washington* 103: 546-553.
- Borkent, A. and W.L. Grogan. 1995. A revision of the genus *Ceratopogon* Meigen. *Memoirs of the Entomological Society of Washington* 15: 1-198.
- Bousquet, Y. (Ed.). 1991. Checklist of Beetles of Canada and Alaska. Research Branch, Agriculture Canada Publication 1861/E. 430 pp.

- Bradley, T.J. and J. Phillips. 1977. Regulation of rectal secretion in saline-water mosquito larvae living in waters of diverse ionic composition. *Journal of Experimental Biology* 66: 83-96.
- Caira, J.N. 1981. Parasitism of Trichoptera by *Bunodera mediovitellata* (Digenea: Allocreadiidae) and the encapsulation responses. M.Sc. thesis, University of British Columbia, Vancouver.
- Caira, J.N. and G.G.E. Scudder. 1985. *Pseudoalloeocreadium alloneotenicum* (Wootton, 1957) Yamaguti, 1971 (Digenea: Allocreadiidae) from *Psychoglypha alascensis* Banks (Trichoptera: Limnephilidae) in Vancouver, British Columbia. *Journal of Parasitology* 7: 524.
- Caira, J.N. and G.G.E. Scudder. 1987. Hemocyte encapsulation of Epon implants in larvae of *Psychoglypha alascensis* (Trichoptera: Limnephilidae). *Transactions of the American Microscopical Society* 106: 120-125.
- Cannings, R.A. 1975a. A new species of *Chironomus* (Diptera: Chironomidae) from saline lakes in British Columbia. *The Canadian Entomologist* 107: 447-450.
- Cannings, R.A. 1975b. Some Chironomidae (Diptera) new to British Columbia and Canada. *Journal of the Entomological Society of British Columbia* 72: 23-25.
- Cannings, R.A. 1978. The distribution of *Tanypteryx hageni* Selys (Odonata: Petaluridae) in British Columbia. *Journal of the Entomological Society of British Columbia* 75: 18-19.
- Cannings, R.A. 1980. Ecological notes on *Sympetrum madidum* (Hagen) in British Columbia, Canada (Anisoptera: Libellulidae). *Notulae odonatologicae* 1(6): 97-99.
- Cannings, R.A. 1981. The larva of *Sympetrum madidum* (Hagen) (Odonata: Libellulidae). *Pan-Pacific Entomologist* 57(2): 341-346.
- Cannings, R.A. 1982a. Notes on the biology of *Aeshna sitchensis* Hagen (Anisoptera: Aeshnidae). *Odonatologica* 11(3): 219-223.
- Cannings, R.A. 1982b. The larvae of the *Tarnetrum* subgenus of *Sympetrum* with a description of the larva of *Sympetrum nigrocreatum* Calvert (Odonata: Libellulidae). *Advances in Odonatology* 1: 9-14.
- Cannings, R.A. 1988. *Pantala hymenaea* (Say) (Anisoptera: Libellulidae) new to British Columbia, Canada, with notes on its status in the northwestern United States. *Notulae odonatologicae* 3(2): 31-32.
- Cannings, R.A. 1996. The Blue Darners: dragonflies of the genus *Aeshna* in British Columbia. *Cordillera* 3 (1): 28-38.
- Cannings, R.A. 1997. *Tramea lacerata* (Hagen) new to British Columbia, Canada, with notes on its status in the northwestern United States (Anisoptera: Libellulidae). *Notulae odonatologicae* 4: 148-149.
- Cannings, R.A. 2001. Rare dragonflies of British Columbia. *Wildlife in British Columbia at Risk* brochure. British Columbia Ministry of Environment, Lands and Parks, Victoria. 6pp.
- Cannings, R.A. and S.G. Cannings. 1983. The Odonata of the Brooks Peninsula, Vancouver Island, British Columbia. *Journal of the Entomological Society of British Columbia* 80: 46-51.
- Cannings, R.A. and S.G. Cannings. 1987. The Odonata of some saline lakes in British Columbia, Canada: ecological distribution and biogeography. *Advances in Odonatology* 3: 7-22.
- Cannings, R.A. and G.P. Doerksen. 1979. The larva of *Ischnura erratica* Calvert (Odonata: Coenagriidae) with ecological notes on the species. *The Canadian Entomologist* 111(3): 327-331.
- Cannings, R.A. and G.G.E. Scudder. 1978. The littoral Chironomidae (Diptera) of saline lakes in central British Columbia. *Canadian Journal of Zoology* 56: 114-115.
- Cannings, R.A. and G.G.E. Scudder. 1979. Phenology of *Chironomus* spp. in saline lakes of central British Columbia. *Internationale Vereinigung für Theoretische und Angewandte Limnologie Verhandlungen* 20: 2641-2646.
- Cannings, R.A. and K.M. Stuart. 1977. The dragonflies of British Columbia. *British Columbia Provincial Museum Handbook No. 35*, Victoria. 254pp.
- Cannings, R.A., S.G. Cannings and R.J. Cannings. 1980. The distribution of the genus *Lestes* in a saline lake series in central British Columbia, Canada (Zygoptera: Lestidae). *Odonatologica* 9(1): 19-28.
- Cannings, R.A., S.G. Cannings and L.R. Ramsay. 2000. The dragonflies (Insecta: Odonata) of the Columbia Basin, British Columbia: field surveys, collections development and public education. Unpublished report, Royal BC Museum and British Columbia Conservation Data Centre, Victoria. On the Internet at http://livinglandscapes.bc.ca/www_dragon/toc.html.
- Cannings, R.A., S.G. Cannings and L.R. Ramsay. 2001. The dragonflies (Insecta: Odonata) of Northern British Columbia: field surveys, collections development and public education. Unpublished report, Royal B.C. Museum and British Columbia Conservation Data Centre, Victoria.
- Cannings, S.G. 1978. Successful rearing of *Cenocorixa expleta* (Hemiptera: Corixidae) in fresh water. *The Canadian Entomologist* 110: 559.
- Cannings, S.G. 1980. New distributional records of Odonata from northwestern British Columbia. *Syesis* 19: 13-15.
- Cannings, S.G. 1981. The influence of salinity on the cuticular permeability of *Cenocorixa bifida hungerfordi* Lansbury (Hemiptera: Corixidae). *Canadian Journal of Zoology* 59: 1505-1509.

- Cannings, S.G. 1989. New records of slender winter stoneflies (Plecoptera: Capniidae) in British Columbia [Canada]. *Journal of the Entomological Society of British Columbia* 86: 77-80.
- Cannings, S.G. and R.A. Cannings. 1980. The larva of *Coenagrion interrogatum* (Odonata: Coenagrionidae), with notes on the species in the Yukon. *The Canadian Entomologist* 112: 438-441.
- Cannings, S.G. and R.A. Cannings. 1994. The Odonata of northern Cordilleran peatlands. *Memoirs of the Entomological Society of Canada* 169: 89-110.
- Cannings, S.G. and R.A. Cannings. 1997. Dragonflies (Odonata) of the Yukon. pp. 169-200 *In* H.V. Danks and J.A. Downes (Eds.). *Insects of the Yukon. Biological Survey of Canada (Terrestrial Arthropods)*, Ottawa. 1034 pp.
- Cannings, S.G., P.D. Cooper and G.G.E. Scudder. 1988. Effect of temperature on rate of water loss in some waterboatmen (Hemiptera: Corixidae). *Canadian Journal of Zoology* 66: 1877-1883.
- Cannings, S.G., R.A. Cannings, and R.J. Cannings. 1991. The distribution of the dragonflies (Insecta: Odonata) of the Yukon Territory, Canada, with notes on ecology and behaviour. *Royal British Columbia Museum Contributions to Natural Science* 13: 1-26.
- Clemens, W.A., D.S. Rawson and J.L. McHugh. 1939. A biological survey of Okanagan Lake, British Columbia. *Bulletin of the Fisheries Research Board of Canada* 56: 1-70.
- Cooper, P.D., G.G.E. Scudder and G.A. Quamme. 1987. Ion and CO₂ regulation in the freshwater water boatman, *Cenocorixa blaisdelli* (Hung.) (Hemiptera, Corixidae). *Physiological Zoology* 60: 465-471.
- Cooper, P.D., G.G.E. Scudder and G.A. Quamme. 1988. Changes in fluid and ion secretion following stimulation in isolated Malpighian tubules of the water boatman, *Cenocorixa blaisdelli* (Hung.) (Hemiptera, Corixidae). *Journal of Insect Physiology* 34: 79-83.
- Cooper, P.D., G.G.E. Scudder and G.A. Quamme. 1989. Segmental differences in secretion by the Malpighian tubules of the freshwater dwelling corixid, *Cenocorixa blaisdelli* (Hung.) (Corixidae, Hemiptera). *Journal of Insect Physiology* 35: 531-536.
- Corkum, L.D. and D.C. Currie. 1987. Distributional patterns of immature Simuliidae (Diptera) in northwestern North America. *Freshwater Biology* 17: 201-221.
- Costello, R.A. 1982. Light trap collections of *Culicoides* (Diptera: Ceratopogonidae) in the Fraser Valley of British Columbia. *Journal of the Entomological Society of British Columbia* 79: 66-70.
- Courtney, G.W. 1990. Revision of Nearctic mountain midges (Diptera: Deuterophlebiidae). *Journal of Natural History* 24: 81-118.
- Currie, D.C. 1986. An annotated list of and keys to the immature black flies of Alberta (Diptera: Simuliidae). *Memoirs of the Entomological Society of Canada* 134: 1-90.
- Currie, D.C. 1997. Black flies (Diptera: Simuliidae) of the Yukon, with reference to the black-fly fauna of northwestern North America. pp. 563-614 *In* H.V. Danks and J.A. Downes (Eds.). *Insects of the Yukon. Biological Survey of Canada (Terrestrial Arthropods)*, Ottawa. 1034 pp.
- Currie, D.C. and P.H. Adler. 1986. Blackflies (Diptera: Simuliidae) of the Queen Charlotte Islands, British Columbia, with discussion of their origin and description of *Simulium* (*Hellichiella*) *nebulosum* n. sp. *Canadian Journal of Zoology* 64: 218-227.
- Currie, D.C. and I.R. Walker. 1992. Recognition and Palaeohydrologic significance of fossil black fly larvae, with a key to the Nearctic genera (Diptera: Simuliidae). *Journal of Paleolimnology* 6: 37-54.
- Curtis, L.C. 1954. Observations on a black fly pest of cattle in British Columbia (Diptera: Simuliidae). *Proceedings of the Entomological Society of British Columbia* 51: 3-6.
- Curtis, L.C. 1967. The Mosquitoes of British Columbia. *British Columbia Provincial Museum Occasional Paper* 15, Victoria. 90 pp.
- Doerksen, G.P. 1980. Notes on the reproductive behaviour of *Enallagma cyathigerum* (Charpentier) (Zygoptera: Coenagrionidae). *Odonatologica* 9: 293-296.
- Donald, D.B. and R.S. Anderson. 1980. The lentic stoneflies (Plecoptera) from the Continental Divide region of southwestern Canada. *The Canadian Entomologist* 112: 753-758.
- Donald, D.B. and D.E. Patriquin. 1983. The wing length of lentic Capniidae (Plecoptera) and its relationship to elevation and Wisconsin glaciation. *The Canadian Entomologist* 115: 921-926.
- Duckitt, G. 1986. Effects of the mosquito larvicide *Bacillus thuringiensis* on caddis flies. M.P.M. thesis, Simon Fraser University, Burnaby.
- Dunkle, S.W. 2000. *Dragonflies through binoculars*. Oxford University Press, New York. 266 pp.
- Edmunds, G.F., S.L. Jensen and L. Berner. 1976. *The mayflies of North and Central America*. University of Minnesota Press, Minneapolis. 330 pp.
- Efford, I.E. and K. Tsumura. 1973. A comparison of the food of salamanders and fish in Marion Lake, British Columbia. *Transactions of the American Fisheries Society* 102: 33-47.
- Ellis, R.A. and J.H. Borden. 1969. Effect of temperature and other environmental factors on *Notonecta undulata* Say (Hemiptera: Notonectidae). *Pan-Pacific Entomologist* 45: 20-25.

- Filmer, N.J. 1964. A Contribution to the Ephemeroptera of British Columbia. M.Sc. thesis, University of British Columbia, Vancouver.
- Gillespie, B.I. and P. Belton. 1980. Oviposition of *Culex pipiens* in water at different temperatures. Journal of the Entomological Society of British Columbia 77: 34-36.
- Hamilton, A.L. 1965. An analysis of a freshwater benthic community with special reference to the Chironomidae. Ph.D. thesis (in 2 parts), University British Columbia, Vancouver.
- Hardy, G.A. 1955. The natural history of the Forbidden Plateau area, Vancouver Island, British Columbia. BC Provincial Museum Natural History and Anthropology Report 1954: B24-B63.
- Harper, P.P. and K.W. Stewart. 1984. Plecoptera. Pages 182-230 in R. W. Merritt and K. W. Cummins (Eds.). An introduction to the aquatic insects of North America. 2nd ed. Kendall/Hunt Publishing Company, Dubuque.
- Hatch, M.H. 1949. A century of entomology in the Pacific Northwest. University of Washington Press, Seattle. 43 pp.
- Hatch, M.H. 1953-1971. Beetles of the Pacific Northwest: Parts 1-5. University of Washington Press, Seattle.
- Hearle, E. 1926. The mosquitoes of the lower Fraser Valley, British Columbia, and their control. National Research Council of Canada Report 17, Ottawa. 94 pp.
- Hearle, E. 1927. List of mosquitoes of British Columbia. Proceedings of the Entomological Society of British Columbia 24: 11-19.
- Heinrichs, M.L., I.R. Walker, R.W. Mathewes and R.J. Hebda. 1999. Holocene chironomid-inferred salinity and paleovegetation reconstruction from Kilpoola Lake, British Columbia. Geographie physique et Quaternaire 53(2): 211-221.
- Heise, B. no date. Mayflies (Ephemeroptera) and stoneflies (Plecoptera) as indicators of logging effects on stream invertebrates (interim report). <http://www.cariboo.bc.ca/SAC/Heise.htm>. Accessed Nov. 9, 2001.
- Hossack, K. and R.A. Costello. 1979. Predation by *Anisogammarus confervicolus* (Amphipoda: Gammaridea) on *Aedes togoi* (Diptera: Culicidae). Journal of the Entomological Society of British Columbia 76: 20-21.
- Hunter, F.F. and V. Connolly. 1986. A cytotoxic investigation of seven species in the *Eusimulium vernum* group (Diptera: Simuliidae). Canadian Journal of Zoology 64: 296-331.
- Ishii, T. and P. Belton. 1984. Evidence for autogenous egg development in *Culex pipiens* in British Columbia. Journal of the American Mosquito Control Association 44: 598-601.
- Jamieson, G.S. and G.G.E. Scudder. 1977. Food consumption in *Gerris* (Hemiptera). Oecologia (Berlin) 30: 23-41.
- Jamieson, G.S. and G.G.E. Scudder. 1979. Predation in *Gerris* (Hemiptera): reactive distances and locomotion rates. Oecologia (Berlin) 44: 13-20.
- Jansson, A. 1972. Mechanisms of sound production and morphology of the stridulatory apparatus in the genus *Cenocorixa* (Hemiptera: Corixidae). Annales Zoologici Fennici 9: 120-129.
- Jansson, A. 1973. Stridulation and its significance in the genus *Cenocorixa* (Hemiptera: Corixidae). Behaviour 46:1-36.
- Jansson, A. 1974a. Annual periodicity of male stridulation in the genus *Cenocorixa* (Hemiptera: Corixidae). Freshwater Biology 4: 93-98.
- Jansson, A. 1974b. Effect of temperature on stridulatory signals of *Cenocorixa* (Hemiptera: Corixidae). Annales Zoologici Fennici 11: 288-296.
- Jansson, A. and G.G.E. Scudder. 1972. Corixidae (Hemiptera) as predators: rearing on frozen brine shrimp. Journal of the Entomological Society of British Columbia 69: 44-45.
- Jansson, A. and G.G.E. Scudder. 1974. The life cycle and sexual development of *Cenocorixa* species (Hemiptera, Corixidae) in the Pacific Northwest of North America. Freshwater Biology 4: 73-92.
- Jaral, M.S. and G.G.E. Scudder. 1970. The morphology and ultrastructure of the Malpighian tubules and hindgut in *Cenocorixa bifida* (Hung.) (Hemiptera: Corixidae). Zeitschrift für Morphologie und Ökologie der Tiere 68: 269-299.
- Jaral, M.S. and G.G.E. Scudder. 1971. Neurosecretion and water balance in *Cenocorixa bifida* (Hung.) (Hemiptera: Corixidae). Canadian Journal of Zoology 49: 1369-1375.
- Jaral, M.S., G.G.E. Scudder and S. Teraguchi. 1969. Observations on the labium of Corixidae (Hemiptera). Canadian Journal of Zoology 47: 713-715.
- Jewett, S.G. 1959. The stoneflies (Plecoptera) of the Pacific Northwest. Oregon State Monographs 3: 1-95.
- Kangasniemi, B.J. and D.R. Oliver. 1983. Chironomidae (Diptera) associated with *Myriophyllum spicatum* in Okanagan Lake Valley lakes, British Columbia. The Canadian Entomologist 115: 1545-1546.
- Kavanaugh, D.H. and P.H. Arnaud, Jr. 1981. Hugh Bosdin Leech - A Curator's Curator. Pan-Pacific Entomologist 57: 2-42.

- Keen, J.H. 1895. List of Coleoptera collected at Masset, Queen Charlotte Islands, BC. The Canadian Entomologist 27: 165-172, 217-220.
- Keen, J.H. 1905. Beetles from northern British Columbia. The Canadian Entomologist 37: 297-298.
- Kenner, R.D. 1996. A new dragonfly for the Lower Mainland. Vancouver Natural History Society Discovery 25: 143-144.
- Kenner, R.D. 2000a. *Somatochlora kennedyi* (Odonata: Corduliidae): a new species for British Columbia, with notes on geographic variation in size and wing venation. Journal of the Entomological Society of British Columbia 97: 47-49.
- Kenner, R.D. 2000b. Lower Mainland dragonfly records for 1999. Vancouver Natural History Society Discovery 29: 22-23.
- Kenner, R.D. 2000c. Annotated checklist of dragonflies of the Lower Mainland. Vancouver Natural History Society Discovery 29: 53-59.
- Kenner, R.D. 2000d. *Gyrinus cavatus* and *G. minutus* (Coleoptera: Gyrinidae) in British Columbia with comments on their nearctic distributions. Journal of the Entomological Society of British Columbia 97: 67-71.
- Kenner, R.D. 2001. Redescription of the larva of *Leucorrhinia glacialis* Hagen with a key to the nearctic *Leucorrhinia* species (Anisoptera: Libellulidae). Odonatologica 30: 281-288.
- Kenner, R.D. and I. Lane. 1997. Photographic records for Lower Mainland dragonflies. Vancouver Natural History Society Discovery 26: 51-53.
- Kenner, R.D. and R.A. Cannings. 2001. Rediscovery of the Western Pondhawk (*Erythemis collocata* [Hagen]) in the Lower Fraser Valley. Vancouver Natural History Society Discovery 29: 50-52.
- Kenner, R.D., R.A. Cannings and S.G. Cannings. 2000. The larva of *Leucorrhinia patricia* Walker (Odonata: Libellulidae). International Journal of Odonatology 3(1): 1-10.
- Kiceniuk, J. and J. Phillips. 1974. Magnesium regulation in mosquito larvae (*Aedes campestris*) living in waters of high MgSO₄ content. Journal of Experimental Biology 61: 749-760.
- Lancaster, J. and G.G.E. Scudder. 1987. Aquatic Coleoptera and Hemiptera in some Canadian saline lakes: patterns in community structure. Canadian Journal of Zoology 65: 1383-1390.
- Lane, I. 2000. First breeding record for the Black Petaltail dragonfly in Canada. Vancouver Natural History Society Discovery 29: 20-21.
- Larson, D.J., Y. Alarie, and R.E. Roughley. 2000. Predaceous Diving Beetles (Coleoptera: Dytiscidae) of the Nearctic Region, with emphasis on the fauna of Canada and Alaska. NRC Research Press, Ottawa. 982 pp.
- Leech, H.B. 1937. A new North American *Agabus*, with notes on other species (Coleoptera: Dytiscidae). The Canadian Entomologist 69: 146-150.
- Leonhardt, K.G. 1985. A cytological study of species in the *Eusimulium aureum* group (Diptera: Simuliidae). Canadian Journal of Zoology 63: 2043-2061.
- Lo, S.E. and A.B. Acton. 1969. The ultrastructure of the rostral sensory organs of the water bug, *Cenocorixa bifida* (Hungerford) (Hemiptera). Canadian Journal of Zoology 47: 717-722.
- Madahar, D.P. 1969. The salivary gland chromosomes of seven taxa in the subgenus *Stegopterna* (Diptera, Simuliidae, *Cnephia*). Canadian Journal of Zoology 47: 115-119.
- Maddrell, S.P.H. and J. Phillips. 1975. Regulation and active transport of sulphate ions by the Malpighian tubules of the saltwater mosquito larvae, *Aedes campestris*. Journal of Experimental Biology 62: 202-216.
- Mahrt, J.L. 1982. Blackflies from Hardwicke Island, British Columbia. Canadian Journal of Zoology 60: 3364-3369.
- Maw, H.E.L., R.G. Footitt, K.G.A. Hamilton and G.G.E. Scudder. 2000. Checklist of the Hemiptera of Canada and Alaska. NRC Research Press, Ottawa. 220 pp.
- Maynard, K.J. 1969. A population study of water-striders (Gerridae: Hemiptera) in Marion Lake, BC. M.Sc. thesis, University of British Columbia, Vancouver.
- McAlpine, J.F. (Ed.). 1981. Manual of Nearctic Diptera, Volume 1. Research Branch, Agriculture Canada, Ottawa.
- McAlpine, J.F. (Ed.). 1987. Manual of Nearctic Diptera, Volume 2. Research Branch, Agriculture Canada, Ottawa.
- McCafferty, W.P. and R.P. Randolph. 1998. Canada mayflies: a faunistic compendium. Proceedings of the Entomological Society of Ontario 129: 47-97.
- McCafferty, W.P., M.J. Wigle and R.D. Waltz. 1994. Systematics and biology of *Acentrella turbida* (McDunnough) (Ephemeroptera: Baetidae). Pan-Pacific Entomologist 70(4): 301-308.
- McMullen, R.D. 1978. Culicoides (Diptera: Ceratopogonidae) of the Okanagan area of British Columbia. The Canadian Entomologist 110: 1053-1057.

- Meredith, J. and J. Phillips. 1973a. Rectal ultrastructure in salt and freshwater mosquito larvae in relation to physiological state. *Zeitschrift für Zellforschung und Mikroskopische Anatomie* 138: 1-22.
- Meredith, J. and J. Phillips. 1973b. The ultrastructure of the anal papillae of a saltwater mosquito larvae *Aedes campestris*. *Journal of Insect Physiology* 19: 1157-1172.
- Meredith, J. and J. Phillips. 1973c. The ultrastructure of anal papillae from a saltwater mosquito larva, *Aedes* (Finlaya) *togoi*. *Canadian Journal of Zoology* 51: 349-353.
- Merritt, R.W. and K.W. Cummins. 1996. An introduction to the aquatic insects of North America. 3rd ed. Kendall/Hunt Publishing Company, Dubuque. 862 pp.
- Minchella, D.J. and M.E. Scott. 1991. Parasitism: A cryptic determinant of animal community structure. *Trends in Ecology and Evolution* 6: 250-254.
- Morley, R.L. and R.A. Ring. 1972a. The intertidal Chironomidae of British Columbia. 1. Keys to their life stages. *The Canadian Entomologist* 104: 1093-1098.
- Morley, R.L. and R.A. Ring. 1972b. The intertidal Chironomidae of British Columbia. 2. Life history and population dynamics. *The Canadian Entomologist* 104: 1099-1121.
- Muchow, C.L. and J.S. Richardson. 2000. Unexplored diversity: macroinvertebrates in coastal British Columbia headwater streams. pp. 503-506 (in) Darling, L.M. (Ed.). *Proceedings of a Conference on the Biology and Management of Species and Habitats at Risk*. Kamloops, B.C. 5-19 Feb. 1999. Volume 2. B.C. Ministry of Environment, Lands and Parks, Victoria, and University College of the Cariboo, Kamloops.
- Mundie, J.H. 1971. The diel drift of Chironomidae in an artificial stream and its relation to diet of Coho salmon fry, *Oncorhynchus kisutch* (Waulbaum). *The Canadian Entomologist* 102: 289-299.
- Munroe, E.G. 1951. The identity and genetic position of *Chauliodes disjunctus* Walker (Megaloptera: Corydalidae). *The Canadian Entomologist* 83: 33-35.
- Munroe, E.G. 1953. *Chauliodes disjunctus*: a correction, with descriptions of a new species and a new genus (Megaloptera: Corydalidae). *The Canadian Entomologist* 85: 190-192.
- Needham, J.G., M.J. Westfall, Jr., & M.L. May. 2000. *Dragonflies of North America*. Scientific Publishers, Gainesville. 939 pp.
- Needham, K.M. 1990. Specific and seasonal variation in survival and sodium balance at low pH in five species of waterboatmen (Hemiptera: Corixidae). M.Sc. thesis, University of British Columbia, Vancouver.
- Needham, K.M. 1996. An identification guide to the nymphal mayflies (Order Ephemeroptera) of British Columbia. Resource Inventory Committee, BC Ministry of Environment, Lands, and Parks, Victoria. 77 pp.
- Nelson, C.R. and R.W. Baumann. 1989. Systematics and distribution of the winter stonefly genus *Capnia* (Plecoptera: Capniidae) in North America. *Great Basin Naturalist* 49: 289-363.
- Ng, K. and J. Phillips. 1985. Magnesium transport across an insect epithelium: role of Na⁺. *Federation Proceedings* 44(3): 851.
- Nimmo, A.P. 1971. The adult Rhyacophilidae and Limnephilidae (Trichoptera) of Alberta and eastern British Columbia, and their post-glacial origin. *Quaestiones Entomologicae* 7: 3-234.
- Nimmo, A.P. 1974. The adult Trichoptera (Insecta) of Alberta and eastern British Columbia, and their post-glacial origins. II. The families Glossosomatidae and Philopotamidae. *Quaestiones Entomologicae* 10: 315-349.
- Nimmo, A.P. 1977. The adult Trichoptera (Insecta) of Alberta and eastern British Columbia, and their post-glacial origins. I. The families Rhyacophilidae and Limnephilidae. Supplement 1. *Quaestiones Entomologicae* 13: 25-67.
- Nimmo, A.P. and G.G.E. Scudder. 1978. An annotated checklist of the Trichoptera (Insecta) of British Columbia. *Syesis* 11: 117-133.
- Nimmo, A.P. and G.G.E. Scudder. 1983. Supplement to an annotated checklist of the Trichoptera (Insecta) of British Columbia. *Syesis* 16: 71-83.
- Northcote, T.G. 1964. Use of a high-frequency echo sounder to record distribution and migration of *Chaoborus* larvae. *Limnology and Oceanography* 9: 87-91.
- Northcote, T.G. and P.A. Larkin. 1963. Western Canada. pp. 451-485. In: *Limnology of North America*, D.C. Frey (Ed.). University of Wisconsin Press, Madison.
- Oliver, D.R. 1984. Description of a new species of *Cricotopus* van der Wulp (Diptera: Chironomidae) associated with *Myriophyllum spicatum*. *The Canadian Entomologist* 116: 1287-1292.
- Oliver, D.R. and M.E. Roussel. 1983. The Insects and Arachnids of Canada: Part 11. The genera of larval midges of Canada (Diptera: Chironomidae). Agriculture Canada Publication 1746, Ottawa. 263 pp.
- Oloffs, P.C. and G.G.E. Scudder 1966. The transition phenomenon in relation to the penetration of water through the cuticle of an insect, *Cenocorixa expleta* (Uhler). *Canadian Journal of Zoology* 44: 621-630.

- Palmer, S.L. 1998. Subfossil chironomids (Insecta: Diptera) and climatic change at high elevation lakes in the Engelmann spruce-subalpine fir zone in southwestern British Columbia. M.Sc. thesis, University of British Columbia, Vancouver.
- Parfin, S.I. and A.B. Gurney. 1956. The spongilla-flies, with special reference to those of the Western Hemisphere (Sisyridae, Neuroptera). *Proceedings of the United States National Museum* 105: 421-529.
- Parkinson, D.A. and R.A. Ring. 1983. Osmoregulation and respiration in a marine chironomid larva, *Paraclunio alaskensis* Coquillett (Diptera: Chironomidae). *Canadian Journal of Zoology* 61: 1937-1943.
- Paulson, D.R. 1997. The dragonflies of Washington. *Bulletin of American Odonatology* 4(4): 75-90.
- Paulson, D.R. 1999. Dragonflies of Washington. Seattle Audubon Society, Seattle. 32 pp.
- Paulson, D.R. and R.A. Cannings. 1980. Distribution, natural history and relationships of *Ischnura erratica* Calvert (Zygoptera: Coenagrionidae). *Odonatologica* 9(2): 147-153.
- Pearlstone, P.S.M. 1973. The food of damselfly larvae in Marion Lake, British Columbia. *Syesis* 6: 33-39.
- Peters, G. 1998. Taxonomic and population studies of British Columbia *Aeshna* species. *Bulletin of American Odonatology* 5: 33-42.
- Peterson, B.V. 1970. The *Prosimulium* of Canada and Alaska. *Memoirs of the Entomological Society of Canada* 69: 1-216.
- Phillips, J. and T.J. Bradley. 1978. Osmotic and ionic regulation in saline-water mosquito larvae. pp. 709-734 (in) Gupta, B.L., R.B. Moreton, J.L. Oschman and B.J. Wall (Eds.). *Transport of Ions and Water in Animal Tissues*. Academic Press, London.
- Phillips, J. and S.P.H. Maddrell. 1974. Active transport of magnesium by Malpighian tubules of the salt water mosquito larvae (*Aedes campestris*). *Journal of Experimental Biology* 61: 761-771.
- Phillips, J. and J. Meredith. 1969a. Osmotic and ionic regulation in a saltwater mosquito larva (*Aedes campestris*). *American Zoologist* 9: 588.
- Phillips, J. and J. Meredith. 1969b. Active sodium and chloride transport by anal papillae of a salt water mosquito larva (*Aedes campestris*). *Nature* 222: 168-169.
- Phillips, J., T.J. Bradley and S.P.H. Maddrell. 1978. Mechanisms of ionic and osmotic regulation in saline-water mosquito larvae. pp. 151-171 (in) Schmidt-Nielsen, Bolis and Maddrell (Eds.). *Comparative Physiology – Water, Ions and Fluid Mechanisms*. Cambridge University Press, Cambridge.
- Pritchard, G. 1989. The roles of temperature and diapause in the life history of a temperate-zone dragonfly: *Argia vivida* (Odonata: Coenagrionidae). *Ecological Entomology* 14: 99-108.
- Pritchard, G. 1991. Insects in thermal springs. *Memoirs of the Entomological Society of Canada* 155: 89-106.
- Provonsha, A.V. 1990. A revision of the genus *Caenis* in North America (Ephemeroptera: Caenidae). *Transactions of the American Entomological Society* 116(4): 801-884.
- Ramsay, L. and S.G. Cannings. 2000. Dragonflies at Risk in British Columbia *In* L. Darling (Ed.). *Proceedings of a Conference on the Biology and Management of Species and Habitats At Risk*, Kamloops, BC. 15 - 19 Feb. 1999. Volume One. BC Ministry of Environment, Lands and Parks and University College of the Cariboo. 490 pp.
- Rawson, D.S. 1934. Productivity studies in lakes of the Kamloops region, British Columbia. *Bulletin of the Biology Board of Canada (Fisheries)* 42: 1-31.
- Reece, P.F. and J.S. Richardson. 2000. Benthic macroinvertebrate assemblages of coastal and continental streams and large rivers of southwestern British Columbia, Canada. *Hydrobiologia* 439: 77-89.
- Rempel, L.L., J.S. Richardson and M.C. Healey. 1999. Flow refugia for benthic invertebrates during flooding of a large river. *Journal of the North American Benthological Society* 18: 34-48.
- Rempel, L.L., J.S. Richardson and M.C. Healey. 2000. Macroinvertebrate community structure along gradients of hydraulic and sedimentary conditions in a large gravel-bed river. *Freshwater Biology* 45: 57-73.
- Reynolds, J.D. 1975. Feeding in corixids (Heteroptera) of small alkaline lakes in central BC. *Internationale Vereinigung für Theoretische und Angewandte Limnologie Verhandlungen* 19: 3073-3078.
- Reynolds, J.D. and G.G.E. Scudder. 1987a. Experimental evidence of the fundamental feeding niche in *Cenocorixa* (Hemiptera: Corixidae). *Canadian Journal of Zoology* 65: 967-973.
- Reynolds, J.D. and G.G.E. Scudder. 1987b. Serological evidence of realized feeding niche in *Cenocorixa* species (Hemiptera: Corixidae) in sympatry and allopatry. *Canadian Journal of Zoology* 65: 974-980.
- Richardson, J.S. 1991. Seasonal food limitation of detritivores in a Montane stream: an experimental test. *Ecology* 72: 873-887.
- Richardson, J.S. 2002. Life cycle phenology of common detritivores from a temperate rainforest stream. *Hydrobiologia*. In press

- Richardson, J.S. and P.M. Kiffney. 2000. Responses of a macroinvertebrate community from a pristine southern British Columbia, Canada, stream to metals in experimental mesocosms. *Environmental Toxicology and Chemistry* 19: 736-743.
- Ricker, W.E. 1943. Stoneflies of Southwest British Columbia. Indiana University Press, Bloomington. 145 pp.
- Ricker, W.E. 1954. Nomenclatorial notes on Plecoptera. *Proceedings of the Entomological Society of British Columbia* 51: 37-39.
- Ricker, W.E. and G.G.E. Scudder. 1976. An annotated checklist of the Plecoptera (Insecta) of British Columbia. *Syesis* 8: 333-348.
- Rosenberg, D.M. and H.V. Danks (Eds.). 1987. Aquatic insects of peatlands and marshes in Canada. *Memoirs of the Entomological Society of Canada* 140: 1-174.
- Ross, H.H. 1937. Studies of Nearctic Aquatic Insects: 1. Nearctic alder flies of the genus *Sialis* (Megaloptera, Sialidae). *Bulletin of the Illinois Natural History Survey* 21: 57-78.
- Ross, H.H. and G.J. Spencer. 1952. A preliminary list of the Trichoptera of British Columbia. *Proceedings of the Entomological Society of British Columbia* 48: 43-51.
- Rowe, L. 1992. Convenience polyandry in a water strider: foraging conflicts and female control of copulation frequency and guarding duration. *Animal Behavior* 44: 189-202.
- Rowe, L. 1994. The costs of mating and mate choice in water striders. *Animal Behaviour* 48: 1049-1056.
- Rowe, L. and G.G.E. Scudder. 1990. Reproductive rate and longevity in the waterstrider *Gerris buenoi*. *Canadian Journal of Zoology* 68: 399-402.
- Rück, A., I.R. Walker and R.J. Hebda. 1998. A palaeolimnological study of Tugulnuit Lake, British Columbia, Canada, with special emphasis on river influence as recorded by chironomids in the lake's sediment. *Journal of Paleolimnology* 19: 63-75.
- Saether, O.A. 1970. Nearctic and palaearctic *Chaoborus* (Diptera:Chaoboridae). *Bulletin of the Fisheries Research Board of Canada* 174: 1-57.
- Sargent, R.W. 1978. Salinity and the physiology of three chironomid species which inhabit saline lakes. M.Sc. thesis, University of British Columbia, Vancouver.
- Schmid, F. 1980. Les Insectes et Arachnides du Canada: Partie 7. Genera des Trichoptères du Canada et des États adjacents. Agriculture Canada Publication 1692, Ottawa. 296 pp.
- Schmid, F. 1998. The Insects and Arachnids of Canada: Part 7. Genera of the Trichoptera of Canada and adjoining or adjacent United States. NRC Research Press, Ottawa. 319 pp.
- Schmid, F. and R. Guppy. 1952. An annotated list of Trichoptera collected on southern Vancouver Island. *Proceedings of the Entomological Society of British Columbia* 48: 41-42.
- Schultze, D.C. and T.G. Northcote. 1972. An experimental study of feeding behavior and interaction of coastal cutthroat trout (*Salmo clarki clarki*) and Dolly Varden (*Salvelinus malma*). *Journal of the Fisheries Research Board of Canada* 29: 555-565.
- Scudder, G.G.E. 1964. Wing muscle polymorphism in *Cenocorixa*. *American Zoologist* 4: 248.
- Scudder, G.G.E. 1965a. The distribution and physiology of some lake Corixidae. *Proceedings of the XII International Congress of Entomology*: 410-411.
- Scudder, G.G.E. 1965b. The Notonectidae of British Columbia. *Proceedings of the Entomological Society of British Columbia* 62: 38-41.
- Scudder, G.G.E. 1966a. The immature stages of *Cenocorixa bifida* (Hung.) and *C. expleta* (Uhler) (Hemiptera: Corixidae). *Journal of the Entomological Society of British Columbia* 63: 33-40.
- Scudder, G.G.E. 1966b. *Notonecta borealis* Bueno and Hussey (Hem., Notonectidae): a flightless species? *Entomologist Monthly Magazine* 102: 258-259.
- Scudder, G.G.E. 1969a. The distribution of two species of *Cenocorixa* in inland saline lakes of British Columbia. *Journal of the Entomological Society of British Columbia* 66: 32-41.
- Scudder, G.G.E. 1969b. The fauna of saline lakes on the Fraser Plateau in British Columbia. *Internationale Vereinigung für Theoretische und Angewandte Limnologie Verhandlungen* 17: 430-439.
- Scudder, G.G.E. 1971a. The osmoregulation and distribution of two species of *Cenocorixa* (Hemiptera). *Proceedings of the XIII International Congress of Entomology, Moscow* 1: 555-556.
- Scudder, G.G.E. 1971b. The postembryonic development of the indirect flight muscles in *Cenocorixa bifida* (Hung.) (Hemiptera: Corixidae). *Canadian Journal of Zoology* 49: 1387-1398.
- Scudder, G.G.E. 1971c. The Gerridae (Hemiptera) of British Columbia. *Journal of the Entomological Society of British Columbia* 68: 3-10.
- Scudder, G.G.E. 1975. Field studies on the flight muscle polymorphism in *Cenocorixa* (Hemiptera: Corixidae). *Internationale Vereinigung für Theoretische und Angewandte Limnologie Verhandlungen* 19: 3064-3072.
- Scudder, G.G.E. 1976. An annotated checklist of the Ephemeroptera (Insecta) of British Columbia. *Syesis* 8: 311-315.

- Scudder, G.G.E. 1977. An annotated checklist of the aquatic and semiaquatic Hemiptera (Insecta) of British Columbia. *Syesis* 10: 31-38.
- Scudder, G.G.E. 1983. A review of factors governing the distribution of two closely related corixids in the saline lakes of British Columbia. *Hydrobiologia* 105: 143-154.
- Scudder, G.G.E. 1987. Aquatic and semiaquatic Hemiptera of peatlands and marshes in Canada. *Memoirs of the Entomological Society of Canada* 140: 65-98.
- Scudder, G.G.E. and G.S. Jamieson. 1972. The immature stages of *Gerris* (Hemiptera) in British Columbia. *Journal of the Entomological Society of British Columbia* 69: 72-79.
- Scudder, G.G.E. and J. Meredith. 1972. Temperature-induced development in the indirect flight muscle of adult *Cenocorixa* (Hemiptera: Corixidae). *Developmental Biology* 29: 330-336.
- Scudder, G.G.E., M.S. Jarial and S.K. Choy. 1972. Osmotic and ionic balance in two species of *Cenocorixa* (Hemiptera). *Journal of Insect Physiology* 18: 883-895.
- Scudder, G.G.E., R.A. Cannings and K.M. Stuart. 1976. An annotated checklist of the Odonata (Insecta) of British Columbia. *Syesis* 9: 143-162.
- Shewell, G.E. 1957. Interim report on distributions of the black flies (Simuliidae) obtained in the northern insect survey. Canadian Department of National Defence, Defence Research Board Environmental Protection Technical Report 7, Ottawa. 50 pp.
- Shewell, G.E. 1959. New Canadian Black flies (Diptera: Simuliidae). III. *Canadian Entomologist* 91: 686-697.
- Smetana, A. 1988. Review of the Family Hydrophilidae of Canada and Alaska (Coleoptera). *Memoirs of the Entomological Society of Canada* 142. 316 pp.
- Smith, B.P. 1977. Water mite parasitism of water boatmen (Hemiptera: Corixidae). M.Sc. thesis, University of British Columbia, Vancouver.
- Soluk, D.A. and J.S. Richardson. 1997. The role of stoneflies in enhancing growth of trout: a test of the importance of predator-predator facilitation within a stream community. *Oikos* 80: 214-219.
- Spence, J.R. 1981. Experimental analysis of habitat selection in waterstriders (Heteroptera: Gerridae). *Ecology* 63: 1505-1514.
- Spence, J.R. 1983. Pattern and process in co-existence of water-striders. *Journal of Animal Ecology* 52: 497-511.
- Spence, J.R. 1989. The habitat templet and life-history strategies of pondskaters (Heteroptera: Gerridae): reproductive potential, phenology and diapause. *Canadian Journal of Zoology* 67: 2433-2447.
- Spence, J.R. 1990. Introgressive hybridization in Heteroptera: the example of *Limnporus* Stål (Gerridae) species in western Canada. *Canadian Journal of Zoology* 68: 1770-1782.
- Spence, J.R. and D. R. Maddison. 1986. Chromosomes of two hybridizing species of *Limnporus* (Heteroptera: Gerridae). *Proceedings of the Entomological Society of Washington* 88: 502-508.
- Spence, J.R. and G.G.E. Scudder. 1978. Larval taxonomy and distribution of *Gerris pingreensis* and *G. incognitus* (Hemiptera: Gerridae) in British Columbia. *Journal of the Entomological Society of British Columbia* 75: 41-45.
- Spence, J.R. and G.G.E. Scudder. 1980. Habitats, life cycles, and guild structure among water striders (Heteroptera: Gerridae) on the Fraser Plateau of British Columbia. *The Canadian Entomologist* 112: 779-792.
- Spence, J.R. and R.S. Wilcox. 1986. The mating system of two hybridizing species of water striders (Gerridae). II. Alternative tactics of males and females. *Behavioral Ecology and Sociobiology* 19: 87-95.
- Spence, J.R., D.H. Spence and G.G.E. Scudder. 1980a. The effects of temperature on growth and development of water strider species (Heteroptera: Gerridae) of central British Columbia and implications for species packing. *Canadian Journal of Zoology* 58: 1813-1820.
- Spence, J.R., D.H. Spence, and G.G.E. Scudder. 1980b. Submergence behavior in *Gerris*: underwater basking. *American Midland Naturalist* 103: 385-391.
- Spencer, G.J. 1942. A preliminary list of the Neuroptera of British Columbia. *Proceedings of the Entomological Society of British Columbia* 38: 23-28.
- Spencer, G.J. 1952. The 1951 status of our knowledge of the insects of British Columbia. *Proceedings of the Entomological Society of British Columbia* 48: 36-41.
- Sperling, F.A.H. and J.R. Spence. 1990. Allozyme survey and relationships of *Limnporus* Stål species (Heteroptera: Gerridae). *The Canadian Entomologist* 122: 29-42.
- Sperling, F.A.H. and J.R. Spence. 1991. Structure of an asymmetric hybrid zone between two water strider species (Hemiptera: Gerridae: *Limnporus*). *Evolution* 45: 1370-1383.
- Sperling, F.A.H., J.R. Spence and N.M. Andersen. 1997. Mitochondrial DNA, allozymes, morphology, and hybrid compatibility in *Limnporus* water striders (Heteroptera: Gerridae): do they all track species phylogenies? *Annals of the Entomological Society of America* 90: 401-415.

- Stace-Smith, G. 1929. Coleoptera of Copper Mt., BC. Museum and Art Notes 4: 69-74.
- Stace-Smith, G. 1930. Coleoptera of Copper Mt., BC. Museum and Art Notes 5: 22-25.
- Stanger, J.A. and R.W. Baumann. 1993. A revision of the stonefly genus *Taenionema* (Plecoptera: Taeniopterygidae). Transactions of the American Entomological Society 119: 171-229.
- Stark, B.P. 2001. North American Stonefly List. <http://www.mc.edu/campus/users/stark/Sfly0102.htm>. Updated February 16, 2001. Accessed November 9, 2001.
- Stark, B.P. and C.R. Nelson. 1994. Systematics, phylogeny and zoogeography of the genus *Yoraperla* (Plecoptera: Peltoperlidae). Entomologica Scandinavica 25: 241-273.
- Stark, B.P., S.W. Szczytko and R.W. Baumann. 1986. North American stoneflies (Plecoptera): Systematics, distribution and taxonomic references. Great Basin Naturalist 46: 383-397.
- Stewart, K.W. and B.P. Stark. 1988. Nymphs of North American stonefly genera (Plecoptera). Thomas Say Foundation of the Entomological Society of America, Hyattsville. 460 pp.
- Stone, A., C.W. Sabrosky, W.W. Wirth, R.H. Foote and J.R. Coulson (Eds.). 1965. A catalogue of the Diptera of North America north of Mexico. USDA Agriculture Handbook 276, Washington, DC. 1696 pp.
- Strange, K. and J. Phillips. 1984. Mechanisms of total CO₂ transport in the microperfused rectal salt gland of *Aedes dorsalis*. I. Ionic requirements of total CO₂ secretion. American Journal of Physiology 246: R727-R734.
- Strange, K. and J. Phillips. 1985. Cellular mechanism of HCO₃⁻ and Cl⁻ transport in insect salt gland. Journal of Membrane Biology 83: 25-37.
- Strange, K., J. Phillips and G.A. Quamme. 1982. Active HCO₃⁻ secretion in the rectal salt gland of a mosquito larva inhabiting NaHCO₃-CO₂ lakes. Journal of Experimental Biology 101: 171-186.
- Strange, K., J. Phillips and G.A. Quamme. 1984. Mechanisms of total CO₂ transport in the microperfused rectal salt gland of *Aedes dorsalis*. II. Site of Cl⁻/HCO₃⁻ exchange and function of anterior and posterior salt gland segments. American Journal of Physiology 246: R735-R740.
- Summerbell, R.C. and R.A. Cannings. 1981. Fungi and *Brillia retifinis* (Diptera: Chironomidae) in the decomposition of conifer leaves in a British Columbia stream. Syesis 14: 125-130.
- Szibbo, C.M. and G.G.E. Scudder. 1979. Secretory activity of the segmented Malpighian tubules of *Cenocorixa bifida* (Hung.) (Hemiptera, Corixidae). Journal of Insect Physiology 25: 931-937.
- Teraguchi, M. and T.G. Northcote. 1966. Vertical distribution and migration of *Chaoborus flavicans* in Corbett Lake, British Columbia. Limnology and Oceanography 11: 164-176.
- Topping, M.S. 1969. Giant chromosomes, ecology, and adaptation. Ph.D. thesis, University of British Columbia, Vancouver.
- Topping, M.S. 1971. Ecology of larvae of *Chironomus tentans* (Diptera: Chironomidae) in saline lakes in central British Columbia. The Canadian Entomologist 103: 328-338.
- Topping, M.S. 1972. The distribution of *Chironomus tentans* in some lakes in central British Columbia in relation to some physical and chemical factors. Proceedings of the XIII International Congress of Entomology (Moscow) 3: 477-478.
- Topping, M.S. and A.B. Acton. 1976. Influence of environmental factors on frequencies of certain chromosomal inversions in *Chironomus tentans* Fabricius. Evolution 30: 170-179.
- Walker, E.M. 1912. The North American dragonflies of the genus *Aeshna*. University of Toronto Press, Toronto. 213 pp.
- Walker, E.M. 1925. The North American dragonflies of the genus *Somatochlora*. University of Toronto Studies (Biology) 26: 1-202.
- Walker, E.M. 1953. The Odonata of Canada and Alaska. Volume 1. University of Toronto Press, Toronto. 292 pp.
- Walker, E.M. 1958. The Odonata of Canada and Alaska. Volume 2. University of Toronto Press, Toronto. 318 pp.
- Walker, E.M. and P.S. Corbet. 1975. The Odonata of Canada and Alaska. Volume 3. University of Toronto Press, Toronto. 307 pp.
- Walker, I.R., J.P. Smol, D.R. Engstrom and H.J.B. Birks. 1991. An assessment of Chironomidae as quantitative indicators of past climatic change. Canadian Journal of Fisheries and Aquatic Sciences 48: 975-987.
- Walker, I.R., E. Reavie, S. Palmer and R.N. Nordin. 1993. A palaeoenvironmental assessment of human impact on Wood Lake, Okanagan Valley, British Columbia. Quaternary International 20: 51-70.
- Westfall, M.J., Jr. and M.L. May. 1996. Damselflies of North America. Scientific Publishers, Gainesville. 649 pp.
- Whitehouse, F.C. 1941. British Columbia dragonflies (Odonata), with notes on distribution and habits. American Midland Naturalist 26: 488-557.

- Wiggins, G.B. 1977. Larvae of the North American caddisfly genera (Trichoptera). 1st ed. University of Toronto Press, Toronto. 401 pp.
- Wiggins, G.B. 1996. Larvae of the North American caddisfly genera (Trichoptera). 2nd ed. University of Toronto Press, Toronto. 457 pp.
- Wigle, M.J. and H.V. Thommasen. 1990. Ephemeroptera of the Bella Coola and Owikeno Lake watersheds, British Columbia central coast. Journal of the Entomological Society of British Columbia 87: 7-15.
- Wilcox, R.S. and J.R. Spence. 1986. The mating system of two hybridizing species of water striders (Gerridae). I. Ripple signal functions. Behavioral Ecology and Sociobiology 19: 79-85.
- Winterbourn, M.J. 1971a. The life histories and trophic relationships of the Trichoptera of Marion Lake, British Columbia. Canadian Journal of Zoology 49: 623-635.
- Winterbourn, M.J. 1971b. An ecological study of *Banksiola crotchii* Banks (Trichoptera, Phryganeidae) in Marion Lake, British Columbia. Canadian Journal of Zoology 49: 637-645.
- Wood, D.M., P.T. Dang and R.A. Ellis. 1979. The Mosquitoes of Canada (Diptera: Culicidae). The Insects and Arachnids of Canada. Part 6. Agriculture Canada Publication 1686. 390 pp.
- Zloty, J. 1996. A revision of the nearctic *Ameletus* mayflies based on adult males, with descriptions of seven new species (Ephemeroptera: Ameletidae). The Canadian Entomologist 128: 293-346.
- Zloty, J. and G. Pritchard. 1997. Larvae and adults of *Ameletus* mayflies (Ephemeroptera: Ameletidae) from Alberta. The Canadian Entomologist 129: 251-289.

